



BDG SYNTHESIS

Certificate of Analysis

BDG Synthesis certifies that this reference material meets or exceeds the specifications stated herein.


Neil Beare

Neil Beare, PhD, Director
19 January 2015

Name: 2-Hydrazino-1-methylpyridinium Tosylate (HMP)

CAS Number: none

Structure:



Molecular Weight: $C_6H_{10}N_3 \cdot C_7H_7O_3S = 295.36$

Lot Number: BDG 14194.1

Appearance: Off white, crystalline solid

Corrected Purity: 100.0 % (HPLC) - 2.1 % (water) = 97.9 %

Re-test Date: 19 January 2017

Storage and Handling:

Temperature:	refrigerate for prolonged storage; may be handled and shipped at ambient temperature.
Humidity:	not believed to be hygroscopic; may be handled in normal laboratory atmosphere.
Light:	protect from strong sunlight.
Caution:	only experienced laboratory personnel should handle the material.

Identity and Purity

Proton NMR Spectrum

Identity: the signals are consistent with the proposed structure and in accord with literature where available.

Residual Solvents: no residual solvents are observed.

Impurities: no significant impurities are evident in the spectrum.

Carbon-13 NMR Spectrum

Identity: the signals are consistent with the proposed structure and in accord with literature where available.

High-resolution Mass Spectrum (TOF MS ES+)

Found m/z 124.0870. $C_6H_{10}N_3 [M]^+$ (2-hydrazino-1-methylpyridinium) requires m/z 124.0875. The deviation of 4.0 ppm is within normally accepted limits for the establishment of identity by HRMS.

HPLC

A sharp, slightly tailing peak is observed (100.0 %). The broad peak, with elution time of 58 minutes, is derived from toluene-4-sulfonic acid (confirmed by a spiking experiment) and is omitted from the peak area integration.

Note: in the absence of reference materials for preparing calibration curves, it is assumed that all peaks have the same detector response. Where possible, the conditions of analysis follow a pharmacopeial or literature method, or have been adapted from same.

Elemental Analysis

	Found:	C 52.17, H 5.87, N 14.20 %
$C_6H_{10}N_3 \cdot C_7H_7O_3S \cdot 0.2H_2O$	Requires:	C 52.23, H 5.87, N 14.06 %
$C_6H_{10}N_3 \cdot C_7H_7O_3S$	Requires:	C 52.86, H 5.80, N 14.23 %

The elemental analyses fall slightly outside those expected for anhydrous material; the presence of water is reasonably expected from the method of purification and/or the type of material, and the "best-fit" hydrated molecular formula is given.

Karl-Fischer Analysis

	Found:	H ₂ O 2.1 %
$C_6H_{10}N_3 \cdot C_7H_7O_3S \cdot 0.2H_2O$	Requires:	H ₂ O 1.2 %

Of necessity, only a small sample could be used and only a single or duplicate analysis performed. We are unable to state what the errors in the reported water content are, but recommend that the result be used, as the best available, when determining corrected purity.


The available quantity of custom-synthesised material is always small, and this limits the extent and type of analytical data which can be obtained. This Certificate is presented in descriptive format for use by analytical chemists who are trained in the use of custom-synthesised materials. Custom materials often contain higher levels of residual solvents and/or water, and we urge you to use the corrected purity where needed rather than the raw HPLC purity. This compound is intended for use as an analytical reference material and it is not for human administration. Structures are shown with relative stereochemistry unless otherwise specified.

The re-test date is assigned from experience gained with the material in the laboratory and/or on storage. It is not possible to perform formal storage studies because of the small amount of material available.




Proton NMR Spectrum of 2-Hydrazino-1-methylpyridinium Tosylate (HMP) in Acetonitrile-d₃

BDG SYNTHESIS




Lot Number: BDG 14194.1






Carbon-13 NMR Spectrum of 2-Hydrazino-1-methylpyridinium Tosylate (HMP) in Acetonitrile-d₃

BDG SYNTHESIS




Lot Number: BDG 14194.1



BDG - Analysis of 2-Hydrazino-1-methylpyridinium Tosylate (HMP)

Column : Phenomenex Luna C18(2) 5um 250 x 4.6 mm
 Guard : Phenomenex Security Guard C18 RP 4 x 3 mm
 Mobile Phase : 98.5:1.5 50 mM Potassium diHydrogen Phosphate pH=3.0 : Acetonitrile
 Flow Rate : 1.0 mL/min
 Sample Solvent : Mobile Phase
 Column Temperature : 20C
 Injection Volume : 10 uL
 Detection : UV at 234 nm

Sample Name	BDG 14194.1	Instrument	AnalyticalLC01
Acquisition	19/01/2015, 15:36:08	Method (rev.)	LC10555b (10)
Sequence	BDG_19Jan2015b - Reprocessed	Vial Position	1
Operator	solvation010\cerityadmin	Injection	1 of 1



Area Percent Report

Peak#	RT	Peak Height	Peak Area	Width	Area %
1	6.33 min	390.0968	6457.2724	0.2411 min	100.000 %