

Certificate of Analysis

BDG Synthesis certifies that this reference material meets or exceeds the specifications stated herein.

leil beare

Neil Beare, PhD, Director 1 November 2019

Name:

 $7-Hydroxy cannabidiol-d_{10}\\$

CAS Number:

Structure:



Molecular Weight:	$C_{21}H_{20}D_{10}O_3 = 34$	40.52			
Lot Number:	BDG 11660	BDG 11660			
Appearance:	White, crystalline	e solid			
Purity By HPLC:	99.3 %				
Isotopic Purity:	Under 0.5 % d ₀				
Re-test Date:	1 November 2024				
Storage and Handling:	Temperature:	Freeze (-20°C) for prolonged storage; may be handled and shipped at ambient temperature.			
	Humidity:	not believed to be hygroscopic; may be handled in normal laboratory atmosphere.			
	Light:	protect from strong sunlight.			
	Caution:	only experienced laboratory personnel should handle the material.			

Version 1 (Id1267)

Custom synthesis of analytical reference standards, metabolites, stable isotope labelled compounds

Contract research
BDG Synthesis is a division of B Dent Global Limited

Mailing: BDG Synthesis PO Box 38 627, Wellington Mail Centre, Wellington, New Zealand. Shipping: BDG Synthesis Gracefield Research Centre, Building F, Gracefield Road, Lower Hutt, New Zealand.

Phone: + 64 4 569 0520 Fax: + 64 4 569 0521 info@bdg.co.nz www.bdg.co.nz

1/5

Identity and Purity

Proton NMR Spectrum

Identity: the signals are consistent with the proposed structure and in accord with literature where available. Isotopic Labelling: signals at the sites of deuteration are greatly diminished, compared with the spectrum of unlabelled material, indicating clean deuteration.

Residual Solvents: a trace (under 0.1 % w/w) of pentane is observed. Impurities: no significant impurities are evident in the spectrum.

Carbon-13 NMR Spectrum

Identity: the signals are consistent with the proposed structure and in accord with literature where available. Isotopic Labelling: signals at the sites of deuteration have collapsed to small multiplets compared with the spectrum of unlabelled material, indicating clean deuteration.

High-resolution Mass Spectrum (TOF MS ES+)

Found m/z 341.2899. C₂₁H₂₁D₁₀O₃ [M+H]⁺ requires m/z 341.2901. The deviation of 0.6 ppm is within normally accepted limits for the establishment of identity by HRMS. No signal for d₀ material was seen (detection limit about 0.5 %).

HPLC

A sharp, symmetrical peak is observed (99.3 %). Note: in the absence of reference materials for preparing calibration curves, it is assumed that all peaks have the same detector response. Where possible, the conditions of analysis follow a pharmacopeial or literature method, or have been adapted from same.

Elemental Analysis

	Found:	C 74.21, H 5.85, D 5.84 %
$C_{21}H_{20}D_{10}O_3$	Requires:	C 74.07, H 5.92, D 5.91 %

The elemental analyses fall within generally accepted limits for establishing the molecular formula given. The results may also be taken to imply the absence of significant quantities of water or inorganic salts (which have not been elsewhere tested for because of sample size limitations).

The available quantity of custom-synthesised material is always small, and this limits the extent and type of analytical data which can be obtained. This Certificate is presented in descriptive format for use by analytical chemists who are trained in the use of custom-synthesised materials. Custom materials often contain higher levels of residual solvents and/or water, and we urge you to use the corrected purity where needed rather than the raw HPLC purity. This compound is intended for use as an analytical reference material and it is not for human administration. Structures are shown with relative stereochemistry unless otherwise specified.

The re-test date is assigned from experience gained with the material in the laboratory and/or on storage. It is not possible to perform formal storage studies because of the small amount of material available.













				== = =		
Acq. Operator	:	Bruce Hamilton	Seq. Lir	ne :	4	
Acq. Instrument	:	Instrument 1	Locatio	on :	Vial 60	
Injection Date	:	11/1/2019 12:14:21 PM	Ir	nj:	2	
			Inj Volum	ne :	10 µl	
Acq. Method	:	C:\CHEM32\1\METHODS\2018\LC201	L02B.M			
Last changed	:	11/1/2019 10:11:36 AM by Bruce	e Hamiltor	n		
Analysis Method	:	C:\CHEM32\1\METHODS\2018\LC201	02B.M			
Last changed	:	11/1/2019 6:11:11 PM by Bruce	Hamilton			
		(modified after loading)				
Method Info	:	BDG - Analysis of 7-Hydroxycar	nabidiol-	-d10		
		Column : Phenomenex Luna C18	(2) 5 um	250	x 4.6 mm	
		Guard : Phenomenex SecurityGua	ard C18 4	х З	mm	
		Mobile Phase A : 30:70:0.05 Wa	ater : Met	than	ol : Trifluoroacetic Ac	id
		Mobile Phase C : 10:90:0.05 Wa	ater : Met	than	ol : Trifluoroacetic Ac	id
		Gradient (A:C) : T0=100:0, T15	5=0:100, 1	F22=	0:100, T23=100:0,	
		T27=100:0.				
		Sample Solvent : 30:70 Water :	Methanol	1,	Detection : UV 230 nm,	
	• •	Flow : 1 ml/min., Column Tem	nperature	: 3	5 C, Injection : 10	ul.



łr	ea	Per	cent	Report	

Sorted By		:	Sign	nal	
Multiplier		:	1.00	000	
Dilution		:	1.00	000	
Use Multiplier	&	Dilution	Factor	with	ISTDs

Signal 1: DAD1 A, Sig=230,4 Ref=off

Peak #	RetTime [min]	Туре	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1 2 3 4 5 6	9.564 10.083 15.195 17.733 19.403 21.404	BV R VB X BB BB BB BB BB	0.1484 0.1260 0.1422 0.1328 0.1368 0.1771	1.53412e4 16.14060 20.89562 47.80334 7.54900 12.95815	1620.55884 1.88288 2.25355 5.53477 8.73429e-1 1.05630	99.3180 0.1045 0.1353 0.3095 0.0489 0.0839
Total	s:			1.54466e4	1632.15977	

______ *** End of Report ***