



## BDG SYNTHESIS

### Certificate of Analysis

BDG Synthesis certifies that this reference material meets or exceeds the specifications stated herein.


*Barry Dent*

Barry R. Dent, PhD, Director  
19 April 2008

**Name:** Testosterone-19,19,19-d<sub>3</sub>

**CAS Number:** 69660-28-2

**Structure:**



**Molecular Weight:** C<sub>19</sub>H<sub>25</sub>D<sub>3</sub>O<sub>2</sub> = 291.44

**Lot Number:** BDG 3519.4

**Appearance:** White, crystalline solid

**Purity By HPLC:** 99.7 %

**Isotopic Purity:** Under 0.5 % d<sub>0</sub>

**Re-test Date:** 19 April 2013

**Storage and Handling:**

Temperature:	refrigerate for prolonged storage; may be handled and shipped at ambient temperature.
Humidity:	not believed to be hygroscopic; may be handled in normal laboratory atmosphere.
Light:	protect from strong sunlight.
Caution:	only experienced laboratory personnel should handle the material.

## Identity and Purity

### Proton NMR Spectrum

Identity: the signals are consistent with the proposed structure and in accord with literature where available.

Isotopic Labelling: signals at the site of deuteration are absent, compared with the spectrum of unlabelled material, indicating clean deuteration.

Residual Solvents: no residual solvents are observed.

Impurities: no significant impurities are evident in the spectrum.

### Carbon-13 NMR Spectrum

Identity: the signals are consistent with the proposed structure and in accord with literature where available.

Isotopic Labelling: signals at the site of deuteration have collapsed to small multiplets compared with the spectrum of unlabelled material, indicating clean deuteration.

### High-resolution Mass Spectrum (EI+)

Found  $m/z$  291.2281.  $C_{19}H_{25}D_3O_2$   $[M]^+$  requires  $m/z$  291.2278. The deviation of 1.1 ppm is within normally accepted limits for the establishment of identity by HRMS. No signal for  $d_0$  material was seen (detection limit about 0.5 %).

### HPLC

A somewhat broadened, symmetrical peak is observed (99.7 %). Note: in the absence of reference materials for preparing calibration curves, it is assumed that all peaks have the same detector response. Where possible, the conditions of analysis follow a pharmacopeial or literature method, or have been adapted from same.

### Elemental Analysis

	Found:	C 78.13, H 8.65, D 2.17 %
$C_{19}H_{25}D_3O_2$	Requires:	C 78.30, H 8.65, D 2.07 %

The elemental analyses fall within generally accepted limits for establishing the molecular formula given. The results may also be taken to imply the absence of significant quantities of water or inorganic salts (which have not been elsewhere tested for because of sample size limitations).




The available quantity of custom-synthesised material is always small, and this limits the extent and type of analytical data which can be obtained. This Certificate is presented in descriptive format for use by analytical chemists who are trained in the use of custom-synthesised materials. Custom materials often contain higher levels of residual solvents and/or water, and we urge you to use the corrected purity where needed rather than the raw HPLC purity. This compound is intended for use as an analytical reference material and it is not for human administration. Structures are shown with relative stereochemistry unless otherwise specified.

The re-test date is assigned from experience gained with the material in the laboratory and/or on storage. It is not possible to perform formal storage studies because of the small amount of material available.




BDG SYNTHESIS

Proton NMR Spectrum of Testosterone (top) and Testosterone-19,19,19-d<sub>3</sub> (bottom) in C<sub>6</sub>D<sub>6</sub>




Lot Number: BDG 3519.4








**BDG SYNTHESIS**



Carbon-13 NMR Spectrum of Testosterone (top) and Testosterone-19,19,19-d<sub>3</sub> (bottom) in C<sub>6</sub>D<sub>6</sub>




Lot Number: BDG 3519.4



BDG - Gradient analysis of steroid derivatives.

Column : Phenomenex Luna C18(2) 5um 250 x 4.6 mm  
 Guard : Phenomenex Security Guard C18 RP 4 x 3 mm  
 Mobile Phase A : Water  
 Mobile Phase B : Acetonitrile  
 Gradient (A:B) : T0=50:50, T20=40:60, T25=40:60, T28=50:50, T30=50:50  
 Flow Rate : 1.0 mL/min  
 Sample Solvent : Mobile Phase  
 Column Temperature : 20C  
 Injection Volume : 10 uL  
 Detection : UV at 205 nm

<b>Sample Name</b>	BDG 3519.4	<b>Instrument</b>	AnalyticalLC01
<b>Acquisition</b>	19/04/2008, 17:32:26	<b>Method (rev.)</b>	LC10041c ( 5 )
<b>Sequence</b>	BDG_19Apr2008f - Reprocessed	<b>Vial Position</b>	1
<b>Operator</b>	solvation010\cerityadmin	<b>Injection</b>	1 of 1



Area Percent Report

Peak#	RT	Peak Height	Peak Area	Width	Area %
1	5.54 min	0.3639	7.5812	0.2511 min	0.049 %
2	7.87 min	1.5593	21.8150	0.1972 min	0.142 %
3	8.27 min	1.1115	18.1764	0.2528 min	0.118 %
4	9.10 min	1032.3817	15329.8750	0.2132 min	99.691 %